

# AO Images of Asteroids, Inverting their Lightcurves, and SSA

Jack Drummond <sup>a</sup> and Julian Christou <sup>b,c</sup>

<sup>a</sup>*Starfire Optical Range, Directed Energy Directorate  
Air Force Research Laboratory, Kirtland AFB, NM 87117-5776*

<sup>b</sup>*Division of Astronomical Sciences  
National Science Foundation, Arlington Va 22203*

<sup>c</sup>*On leave from Center for Adaptive Optics  
University of California, Santa Cruz, Ca 95064-1077*

## ABSTRACT

In a program to study asteroids with large telescopes, we have recently obtained images of Main-Belt asteroids with adaptive optics (AO) on the Keck-II 10 meter telescope, the world's largest. Although generally featureless as expected, these images show irregular asteroid outlines, and by following the changing size and orientation we have been able to deduce their full triaxial ellipsoid dimensions and spin axis direction in less than one night. Even before the first such AO images, Kaasalainen [1,2] and his colleagues made attempts to deduce the shape of asteroids from their lightcurves. We compare our AO images to these lightcurve inversion (LCI) models, and show the excellent agreement. Similar techniques can be applied to satellite lightcurves.

## 1. INTRODUCTION

As astronomy has been revolutionized by adaptive optics (AO), so has the field of asteroid lightcurve work with new methods of lightcurve inversions. While asteroid shapes (usually in the form of triaxial ellipsoid axes ratios) and rotational pole locations have been obtained from lightcurves (the variation of brightness as the asteroid rotates in several hours) of these unresolved targets over decades as they move around the sky, only in the last several years have these data been turned into (unscaled) three dimensional models by Kaasalainen and his colleagues [1,2]. By making the simple assumptions that the particular light scattering model is unimportant and the asteroid surface is convex everywhere, they have succeeded in inverting lightcurves to construct LCI (LightCurve Inversion) models that deviate from the simple smooth triaxial ellipsoids used in previous lightcurve work, and AO [3,4] as well.

Recalling that no more than a handful of asteroids have ever been visited by spacecraft, and that all asteroids are smaller than the seeing disk ( $< 1$  second of arc) from Earth, the best images of asteroids have been otherwise obtained with AO on 8-10 m telescopes from the ground, the largest being the twin 10 m Keck telescopes on Mauna Kea. In this paper we compare the models of Kaasalainen to images obtained at Keck, and show both the validity of the models and the utility of treating the AO images as projections of our canonical smooth, featureless triaxial ellipsoids.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>SEP 2008</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2008 to 00-00-2008</b>	
4. TITLE AND SUBTITLE <b>AO Images of Asteroids, Inverting their Lightcurves, and SSA</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Air Force Research Laboratory, Directed Energy Directorate, Starfire Optical Range, Kirtland AFB, NM, 87117</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>2008 Advanced Maui Optical and Space Surveillance Technologies Conference, 16-19 Sep, Maui, HI.</b>					
14. ABSTRACT <b>In a program to study asteroids with large telescopes, we have recently obtained images of Main- Belt asteroids with adaptive optics (AO) on the Keck-II 10 meter telescope, the world's largest. Although generally featureless as expected, these images show irregular asteroid outlines, and by following the changing size and orientation we have been able to deduce their full triaxial ellipsoid dimensions and spin axis direction in less than one night. Even before the first such AO images Kaasalainen [1,2] and his colleagues made attempts to deduce the shape of asteroids from their lightcurves. We compare our AO images to these lightcurve inversion (LCI) models, and show the excellent agreement. Similar techniques can be applied to satellite lightcurves.</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

## 2. KECK AO OBSERVATIONS

We competed for heavily oversubscribed time on Keck-II, and were granted one night in August 2006 to image a few asteroids as they rotated during the evening. Using AO behind the 10 m telescope at a wavelength of  $2.1 \mu\text{m}$ , we were able to derive the pole and triaxial ellipsoid dimensions of four asteroids, as reported in Tables 1-3, from images obtained in this single night. Performing a simple deconvolution, dividing the objects by the Lorentzian PSFs obtained in the fitting process, we can compare either convolved or deconvolved images to Kassalainen models. Figure 1 shows the 10 m telescope images of the asteroids and of check stars, and Fig 2 shows their Fast Fourier Transforms (FFT).

Table 1. Asteroid Ellipsoid Parameter Results

Name	$a$ (km)	$b$ (km)	$c$ (km)	$\theta(^{\circ})$	$N(^{\circ})$	$\psi_0$ (UT)
2 Pallas	$548 \pm 3$	$504 \pm 3$	$459 \pm 15$	$+42 \pm 6$	$2 \pm 1$	$8.71 \pm 0.06$
129 Antigone	$163 \pm 4$	$133 \pm 2$	$90 \pm 28$	$+46 \pm 10$	$231 \pm 3$	$10.25 \pm 0.06$
409 Aspasia	$198 \pm 5$	$172 \pm 3$	$172 \pm 3$	$-34 \pm 3$	$293 \pm 5$	$10.01 \pm 0.12$
704 Interamnia	$349 \pm 4$	$339 \pm 3$	$274 \pm 22$	$-26 \pm 16$	$275 \pm 1$	$6.62 \pm 0.19$


Table 2. Asteroid Ellipsoid Poles

Name	Eq		Pole		Ecl	
	RA( $^{\circ}$ )	Dec( $^{\circ}$ )	Err	Rad ( $^{\circ}$ )	$\lambda(^{\circ})$	$\beta(^{\circ})$
2 Pallas	41	-13		3	34	-27
129 Antigone	222	+39		5	202	+52
409 Aspasia	58	+65		4	72	+43
704 Interamnia	343	+73		3	47	+66


Table 3. Volumes and Mean Diameters

Name	Volume ( $\text{km}^3$ )	$\bar{d}$ (km)	IRAS [5] $d$ (km)
2 Pallas	$6.64(\pm 0.22) \times 10^7$	$502 \pm 5$	$498 \pm 19$
129 Antigone	$1.02(\pm 0.32) \times 10^6$	$125 \pm 13$	124 <sup>†</sup>
409 Aspasia	$3.05(\pm 0.11) \times 10^6$	$180 \pm 2$	$162 \pm 7$
704 Interamnia	$1.70(\pm 0.14) \times 10^7$	$319 \pm 9$	$317 \pm 5$

<sup>†</sup> Not observed by IRAS; diameter is from the IMPS ground-based catalog [5].




**Figure 1.** Images of four asteroids and two stars. The star at upper right has a Strehl of 0.47, and at lower right 0.62. Unlike the images of asteroids to follow, these images are displayed on a square root scale to show the faint structure around the stars.



**Figure 2.** Amplitudes of FFT's of images in Fig 3. To show subtle but important detail, these FFT's are displayed on a log scale. Notice how the high frequency cutoffs are well demarcated in all of the FFT's, outside of which appears only noise. Also notice how the large asteroids Pallas and Interamnia show several ripples or minima while the smaller asteroids show only one or two.

### 3. LIGHTCURVE INVERSION MODELS


Kassalainen and colleagues make their models available on a web site\* and in Figs 3 and 4, we compare deconvolved images of 2 Pallas and 129 Antigone to the models. As expected, the images show no albedo features (asteroids have been painted gray by collisions), but deviations from elliptical outlines are apparent, and are well-captured by Kassalainen's models, although some differences between the LCI models and our images are evident.



**Figure 3.** Deconvolved images of 2 Pallas and models from lightcurve inversions.

---

\*<http://astro.troja.mff.cuni.cz/projects/asteroids3D/web.php>




**Figure 4.** Deconvolved images of 129 Antigone and models from lightcurve inversions.

#### 4. EDGE RECONSTRUCTION

With featureless asteroids, image reconstruction techniques are most productive in finding the edges or outlines of objects. We have developed a method to produce the same kind of LCI 3-D models from the contours of a series of reconstructed images of asteroids. Since asteroids are so hard-edged, any unsophisticated algorithm can be used to produce a one-contour outline, and we use the standard MATLAB contour function on images produced by our simple deconvolution. For each image, we know the asteroid's sub-Earth longitude and latitude from our triaxial ellipsoid model and pole. We then construct the coordinates of the contours for each image produced by MATLAB, keeping in mind that edges are defined not by points located  $90^\circ$  from the direction of the observer, but by points where the surface normal is pointing  $90^\circ$  from the observer. The surface normals for ellipsoids are easily and exactly calculated for every point on the surface. Combining these longitudes and latitudes with the apparent distances from the center to the contour points then gives radius vectors in asteroid coordinates for the contours of a single image. By combining

several such contours at various rotational phases (Fig 5), a 3-D model can be built up. Our 3-D model of Antigone from 10 Keck images in one night is shown in Fig 6.




**Figure 5.** Contours on the projected 3-D model of 129 Antigone reconstructed from edges.


Because only limited views of an asteroid can be obtained in one night, much of the surface, and many outlines, are not visible, leaving large gaps in the 3-D model. We can fill in these radius vectors by using the ones from the triaxial ellipsoid model we obtain from fitting the images. Turning over these radius vectors to MATLAB's *convhulln.m* program then produces a set of tetrahedrons, where one face of each contributes to covering the entire surface of an asteroid. This combination ellipsoid and edge model is also shown in Fig 6.

Figure 7 shows four 3-D models viewed from three orthogonal directions. It is apparent that extrapolating to unseen regions can result in losing the true shape, but this can easily be remedied by performing the same analysis at another opposition, when the asteroid has moved significantly, providing a different viewing aspect.





**Figure 6.** 3-D models of Antigone made from 10 deconvolved images using edges only (top ten) and edges supplemented with ellipsoid solution radius vectors (bottom ten).



**Figure 7.** Four 3-D models of Antigone, all normalized to the same volume. Each is shown at the same sub-Earth longitude and latitude as listed on the left.

## 5. SSA

The techniques developed by Kaasalainen and colleagues are based on unresolved photometric measurements of asteroids observed over decades. Low-earth satellites can go through the same geometry in one pass. Edge reconstruction can be applied to a series of two-dimensional deconvolved AO images, again available on one pass. Application of these two techniques could be profitable in the study of satellites, both LEOs and GEOs, although for the edge reconstruction technique the latter would require observations with 8-10 m telescopes. A satellite pass makes the problem easier in that the observations can be obtained quickly, but the overall problem is much harder because the basic assumptions used for asteroids do not necessarily apply to satellites. Satellites are not smooth featureless triaxial ellipsoids, they may have strong concavities, and scattering laws are important (specular, diffuse, Lambertian, etc.). However, with a different set of simplifying assumptions, the techniques used on asteroids can be a tool for studying barely resolved or unresolved satellites.

## 6. REFERENCES

1. Kaasalainen, M., and Torppa, J. 2001. Optimization Methods for Asteroid Lightcurve Inversion. I. Shape Determination. *Icarus* 153, 24-36.
2. Kaasalainen, M., Torppa, J., and Muinonen, K. 2001. Optimization Methods for Asteroid Lightcurve Inversion. II. The Complete Inverse Problem. *Icarus* 153, 37-51.
3. Drummond, J., and Christou, J. 2008. Triaxial ellipsoid dimensions and rotational poles of seven asteroids from Lick Observatory adaptive optics images, and of Ceres. *Icarus* in press, doi:10.1016/j.icarus.2008.05.009
4. Drummond, J.D. 2000. Measuring Asteroids with Adaptive Optics. In: Ageorges, N. Dainty, C. (Eds.), *Laser Guide Star Adaptive Optics for Astronomy*, Kluwer Academic Publishers, Dordrecht, 243-262.
5. Tedesco, E.F., Veeder, G.J., Fowler, J.W. and Chillemi, J.R. 1992. The IRAS Minor Planet Survey – Final Report (PL-TR-92-2049), Phillips Laboratory, Hanscom Air Force Base, MA.